

# RECOMMENDED REGIMEN FOR EXPOSED PATIENTS 13 YEARS + OLDER

WITH NORMAL RENAL FUNCTION (CREATININE CLEARANCE GREATER/EQUAL TO 60 ML/MIN)

## RECOMMENDED nPEP REGIMEN

**TENOFOVIR DF 300 mg/\*EMTRICITABINE 200 mg fixed dose combination (FDC) (TRUVADA®) PO daily  
with  
RALTEGRAVIR (ISENTRESS®) 400 mg twice daily  
or  
DOLUTEGRAVIR (TIVICAY®) 50 mg once daily**

**Duration of therapy: 28-day course of nPEP is recommended.**

\*Lamivudine 300 mg PO daily may be substituted for emtricitabine. An FDC is available when tenofovir is used with emtricitabine.

## ALTERNATIVE nPEP REGIMEN

**TENOFOVIR DF 300 mg/\*EMTRICITABINE 200 mg fixed dose combination (FDC) (TRUVADA®) once daily  
with  
DARUNAVIR 800 mg (as two, 400 mg tablets) (PREZISTA®) once daily  
and  
RITONAVIR 100 mg once daily**

**Duration of therapy: 28-day course of nPEP is recommended.**

\*Lamivudine 300 mg PO daily may be substituted for emtricitabine. An FDC is available when tenofovir is used with emtricitabine.