

RECOMMENDED REGIMEN FOR EXPOSED PATIENTS 13 YEARS + OLDER

WITH NORMAL RENAL FUNCTION (CREATININE CLEARANCE GREATER/EQUAL TO 60 ML/MIN)

RECOMMENDED nPEP REGIMEN

TENOFOVIR DF 300 mg/*EMTRICITABINE 200 mg fixed dose combination (FDC) (TRUVADA®) PO daily
with
RALTEGRAVIR (ISENTRESS®) 400 mg twice daily
or
DOLUTEGRAVIR (TIVICAY®) 50 mg once daily

Duration of therapy: 28-day course of nPEP is recommended.

*Lamivudine 300 mg PO daily may be substituted for emtricitabine. An FDC is available when tenofovir is used with emtricitabine.

ALTERNATIVE nPEP REGIMEN

TENOFOVIR DF 300 mg/*EMTRICITABINE 200 mg fixed dose combination (FDC) (TRUVADA®) once daily
with
DARUNAVIR 800 mg (as two, 400 mg tablets) (PREZISTA®) once daily
and
RITONAVIR 100 mg once daily

Duration of therapy: 28-day course of nPEP is recommended.

*Lamivudine 300 mg PO daily may be substituted for emtricitabine. An FDC is available when tenofovir is used with emtricitabine.